

Local estrogen biosynthesis in males and females

E Simpson¹, G Rubin¹, C Clyne¹, K Robertson¹, L O'Donnell¹, S Davis² and M Jones¹

¹ Prince Henry's Institute of Medical Research, PO Box 5152, Clayton, Victoria 3168, Australia

² Jean Hailes Foundation, 173 Carinish Rd, Clayton, Victoria 3168, Australia

(Requests for offprints should be addressed to E Simpson)

Abstract

It is now apparent that in men and in postmenopausal women, estrogens have important physiological and pathophysiological roles. However, importantly, these actions are at a local level, namely paracrine, autocrine, and even 'intracrine' rather than endocrine in the classical sense. Thus for example local estrogen biosynthesis in the bones of men plays a hitherto unsuspected role in the maintenance of bone mineralization and in epiphyseal fusion; and in the testes, estrogen is essential for male germ cell development. On the other hand, in postmenopausal women, the mesenchymal cells of the breast are the major source of estrogen responsible for breast cancer development. This realization points to the importance of circulating C₁₉ precursors in the maintenance of adequate estrogen biosynthesis in extragonadal sites and suggests the possibility of new therapies to block estrogen synthesis in a tissue-specific fashion.

Endocrine-Related Cancer (1999) 6 131-137

Introduction

In recent years, our understanding of the role of estrogens in both females and males has expanded greatly. For example, considerable emphasis has been focused on the regulation of extragonadal estrogen biosynthesis, in particular that which occurs in adipose tissue and bone, and its importance in the well-being of the elderly (Simpson *et al.* 1997). The regulation of aromatase expression in normal adipose tissue from various body sites including the breast has been examined as a function of age (Bulun & Simpson 1994), and significant changes in the regulation of the expression which occurs in adipose tissue proximal to a breast tumor (Agarwal *et al.* 1996) have been documented. This has led to the conclusion that tumorous epithelium of the breast, and/or macrophages recruited to the tumor site, produce factors such as prostaglandin E₂ (PGE₂), tumor necrosis factor α (TNF α) and class I cytokines, which regulate aromatase expression in the surrounding mesenchymal cells of the adipose tissue and of the tumor itself (Zhao *et al.* 1995, 1996a,b, Fig. 1).

Since bone is a favourite site for breast cancer metastasis, attention has also focused on aromatase expression in osteosarcoma cell lines, in primary cultures of human fetal osteoblasts, as well as osteoclastic cell lines

such as THP-1 (Shozu *et al.* 1997, Shozu & Simpson 1998). We and others have observed that these cells exhibit high expression of aromatase activity which is regulated primarily by class I cytokines, interleukin-1 β (IL-1 β) and TNF α . These observations, together with the observation of a marked bone phenotype in men with mutations of either the estrogen receptor (ER) (α) or aromatase (Smith *et al.* 1994, Carani *et al.* 1997), have led to the conclusion that local estrogen production in bone cells plays an important role in the maintenance of bone mineralization and the prevention of osteoporosis in men and in women. In an extension of these concepts, we have advanced the hypothesis, also enunciated by Labrie and colleagues (1997a), that in postmenopausal women, and also in men, extragonadal estrogen biosynthesis in a number of sites, including adipose tissue, bone, various sites of the brain, vascular endothelial and smooth muscle cells, plays an important but hitherto largely unrecognized physiologic and pathophysiologic role in a paracrine, autocrine and indeed, intracrine, fashion (Simpson & Davis 1998). The long-term health consequences of estrogen decline after the menopause include bone loss, urogenital aging, increased cardiovascular disease, and probably cognitive impairment culminating in dementia. The incidence and pattern of occurrence of all of these disease processes differ significantly between men and




Figure 1 Proposed regulation of aromatase gene expression in breast adipose tissue from cancer-free individuals and from those with breast cancer. In the former case, expression is stimulated primarily by class I cytokines or TNF α produced locally, in the presence of systemic glucocorticoids. As a consequence, promoter I.4-specific transcripts of aromatase predominate. In the latter case, PGE $_2$ produced by the tumorous epithelium, tumor derived fibroblasts, and/or macrophages recruited to the tumor site is the major factor stimulating aromatase expression, as evidenced by the predominance of promoter II and I.3-specific transcripts of aromatase.

DEX, dexamethasone; OSM, oncostatin M; LIF, leukemia inhibitory factor.

women, and cannot be explained by gender differences in circulating estrogen levels alone. For example, men have plasma estradiol levels in the postmenopausal range throughout their adult years, but rarely develop osteoporosis until very late in life. Hence, our understanding of the peripheral metabolism of precursor steroids in the main estrogen target tissues appears fundamental to ascertaining the mechanisms underlying the development of diseases associated with the decline in circulating estrogen levels after menopause.

Of equal significance is the realization derived from studies of ER α -knock-out (ERKO) (Lubahn *et al.* 1993) and aromatase knock-out (ArKO) mice (Fisher *et al.* 1998), of the role of locally-produced estrogen in the regulation of male reproduction and in particular its role in spermatogenesis. This new understanding of the role of estrogens in the male blurs our definition of male versus female hormones, since, at least at the local level, both androgens and estrogens have important roles to play in both sexes.

Sites of estrogen biosynthesis

While the ovaries are the principal source of systemic estrogen in the premenopausal non-pregnant woman,

other sites of estrogen biosynthesis are present throughout the body and these become the major sources of estrogen beyond menopause. These sites include the mesenchymal cells of the adipose tissue and skin (reviewed in Simpson *et al.* 1997), osteoblasts (Bruch *et al.* 1992) and perhaps osteoclasts (Jacob *et al.* 1995) in bone, possibly vascular endothelial (Bayard *et al.* 1995) and aortic smooth muscle cells (Murakami *et al.* 1998) as well as a number of sites in the brain including the medial preoptic/anterior hypothalamus, the medial basal hypothalamus and the amygdala (Naftolin *et al.* 1975). These extragonadal sites of estrogen biosynthesis possess several fundamental features which differ from those of the ovaries. Principally, these sites are dependent on circulating precursor C $_{19}$ steroids for estrogen biosynthesis. Although these extragonadal tissues have the capacity to convert C $_{19}$ steroids to C $_{18}$ steroids, unlike the ovaries they lack the ability to synthesize C $_{19}$ precursors. Hence, estrogen production in adipose, bone and brain is totally dependent on the availability of circulating C $_{19}$ precursors. Another important feature is that the estrogen synthesized within these compartments, particularly bone, breast and brain, is probably only biologically active at a local tissue level in a paracrine or 'intracrine' fashion (Labrie *et al.* 1997). Thus, the total amount of estrogen synthesized by these

extragonadal sites may be small, but the local tissue concentrations achieved are probably quite high, and exert significant biologic influence locally.

After menopause, the mesenchymal cells of the adipose tissue become the main source of estrogen (Siiteri & MacDonald 1973, Simpson *et al.* 1997). Therefore, in the post-reproductive years, the degree of a woman's estrogenization is mainly determined by the extent of her adiposity. This is of clinical importance, since corpulent women are relatively protected against osteoporosis (Melton 1997) and the incidence of Alzheimer's disease is lower in more corpulent postmenopausal women than in their slimmer counterparts (V Henderson, personal communication). On the down-side, obesity is positively correlated with the risk of breast cancer (Huang *et al.* 1997).

In the case of males, the Leydig cells (Tsai-Morris *et al.* 1985) and other cells of the testes, including germ cells in various stages of differentiation (Nitta *et al.* 1993), produce estrogen which, as mentioned previously, has an important role in spermatogenesis. Nevertheless, it has been estimated that at best the testes can account for 15% of circulating estrogens (Hemsell *et al.* 1974) and hence, in the male, local production of estrogens, both intratesticular and extragonadal, is of physiologic significance throughout adult life. For example, estrogen production in bone appears to be vital for the maintenance of bone mineralization and prevention of osteoporosis. This is supported by studies of men, either with a mutation of the gene encoding the aromatase enzyme (Morishima *et al.* 1995, Carani *et al.* 1997) or a mutation of the ER (Smith *et al.* 1994). These individuals exhibit failure of epiphyseal fusion, osteopenia and delayed bone age. Recently, we have observed that male ArKO mice also exhibit alterations in bone histomorphometry characteristic of undermineralization (Oz *et al.* 1998). In a similar fashion, it is reasonable to speculate that estrogen production in one or more brain sites has an influence on sexual behaviour and, as suggested by recent observational epidemiologic studies, may have a role in the maintenance of cognitive function and the prevention of Alzheimer's disease (reviewed in Yaffe *et al.* 1998). In this context it is appropriate to reconsider why osteoporosis is more common in women than in men, and affects women at a younger age, in terms of fracture incidence. Similarly, one may question why the incidence of Alzheimer's disease is greater among women than among men.

Precursor availability

A key factor in the gender difference in the incidence of these diseases appears to be the availability of precursor C₁₉ steroids for aromatization to estrogens in extragonadal

sites, a concept also advanced by Labrie *et al.* (1998). In postmenopausal women the principal source of C₁₉ steroid production is the adrenal cortex, which elaborates androstenedione, dehydroepiandrosterone (DHEA) and DHEA sulfate (DHEAS). However, the secretion of these steroids and their plasma concentrations decrease markedly with advancing age (Labrie *et al.* 1997b). Moreover, DHEA must first be converted to androstenedione prior to aromatization. Another major step is the reduction of the 17-keto group to 17 β -hydroxyl, catalyzed by 17 β -hydroxysteroid dehydrogenase (HSD) type I, which is essential for formation of the active estrogen, estradiol. The distribution of this enzyme in the various extragonadal sites of aromatization has not yet been fully established, although it is expressed in tumorous breast epithelium (Sasano *et al.* 1996) and in bone (Sasano *et al.* 1997). It should be noted in this context that there is a recent report that 17 β -HSD type III, which converts androstenedione to testosterone, is present in visceral fat (Corbould *et al.* 1998), together with 17-HSD type II.

Interestingly, in the male circulation, the levels of testosterone are at least one order of magnitude greater than those circulating in the plasma of postmenopausal women (10-30 vs 0.5 nmol/l), while the levels of androstenedione are rather similar (~2.5 nmol/l). Since the levels of circulating testosterone in the male are similar to the K_m of aromatase (20-30 nmol/l), it is likely that circulating testosterone can be converted efficiently in extragonadal sites to give rise to local concentrations of estradiol sufficient to transactivate both ERs (α and β) (K_D ~1 nmol/l). Moreover, although testosterone levels in the plasma of men decrease with advancing years, this decrease is small compared with the decrease in the circulating levels of adrenal C₁₉ steroids. Consequently, compared with women, males maintain a high circulating level of the active precursor testosterone throughout life, which is available for conversion to the active estrogen, estradiol, in extragonadal sites. Not only is the level of circulating testosterone in men much greater than that in women, but it is also two orders of magnitude greater than the mean levels of circulating estradiol in postmenopausal women (less than 130 pmol/l) and in men (~25-130 pmol/l). Given that most of this circulating estradiol is probably bound to sex hormone binding globulin, it is unlikely to have a major impact on transactivation of the ER, compared with estrogen produced locally as a consequence of conversion of circulating testosterone. Thus, the uninterrupted sufficiency of circulating testosterone in men throughout life supports the local production of estradiol by aromatization of testosterone in estrogen-dependent tissues, and thus affords ongoing protection against the so-called estrogen deficiency diseases. This appears to be important in terms of protecting the bones of men against mineral loss and may contribute to the




Figure 2 Importance of circulating C₁₉ steroids as precursors for extragonadal estrogen biosynthesis in men (left panel) and postmenopausal women (right panel).
E, estradiol; Δ⁴, Δ⁴-androstenedione; T, testosterone.

maintenance of cognitive function and prevention of Alzheimer's disease in men (Fig. 2).

Currently, there is considerable interest in the use of testosterone as a component of hormone replacement therapy (HRT) for postmenopausal women, but its use is mostly limited to those women who complain of loss of sexual interest and libido. However, there is increasing evidence that postmenopausal testosterone replacement is effective in both the prevention and treatment of osteoporosis (Davis *et al.* 1995, Raisz *et al.* 1995). Thus, the present discussion suggests a broader role for the use of testosterone in HRT, namely as a circulating precursor for local synthesis of estrogen in target tissues where the latter acts in an autocrine and paracrine fashion.

Regulation of aromatase expression in adipose tissue

We have suggested previously (Simpson *et al.* 1997) that aromatase is a marker of the undifferentiated adipose mesenchymal cell phenotype. In support of this, the factors which stimulate expression in adipose tissue of cancer-free individuals are factors which either inhibit or reverse the differentiated phenotype of adipocytes, namely class I cytokines such as IL-6, oncostatin M and IL-11 or else TNF α . Moreover, all of these factors act via promoter I.4 of the aromatase gene and require glucocorticoids as co-stimulators (reviewed in Simpson *et al.* 1997). Adipocyte differentiation is driven by trans-

cription factors such as C/EBP and also PPAR γ (Spiegelman 1998), and while involving the down-regulation of aromatase expression, the differentiation process also involves the upregulation of markers such as lipoprotein lipase, the insulin receptor and GLUT4. These actions are antagonized by TNF α which is also expressed in adipocytes (Hotamisligil *et al.* 1993). Significantly in this context, mice lacking TNF α function are protected from obesity-induced insulin-resistance (Uysal *et al.* 1997). These considerations suggest that factors which stimulate adipocyte differentiation such as ligands of the PPAR γ receptor (e.g. BRL 49653 and 15-deoxy- $\Delta^{12,14}$ -PGJ₂) would inhibit aromatase expression and this has proven to be the case (Fig. 3). They also indicate that individuals with insulin-resistance have higher levels of aromatase expression in their adipose tissue, and therefore are at greater risk of developing breast cancer. While the former has not been shown, there is epidemiologic evidence to support the latter contention (Bruning *et al.* 1992).

Further light on the role of estrogens in adipose tissue metabolism has come from our recent studies employing the ArKO mouse, in which there is a redistribution of adipose tissue with diminished subcutaneous and increased visceral fat deposits (Table 1). To investigate the mechanism whereby the subcutaneous fat deposits are decreased, we examined the actions of estrogen on the differentiation of 3T3 L1 cells, and observed that estradiol

Table 1 Body mass and visceral fat content of female ArKO mice and wild-type littermates

		Wild-type	ArKO
Body mass (g)	10-12 weeks	19.3 ± 0.4	21.9 ± 0.1
	4-5 months	21.7 ± 0.3	27.0 ± 3.3
Gonadal fat mass (mg)	10-12 weeks	162.5 ± 13.2	269.6 ± 86.9
	4-5 months	280.0 ± 84.7	490.0 ± 96.0

mimics the effects of troglitazone in this context. So an important issue which arises is to determine the mechanism whereby estradiol elicits this response and in particular to determine whether estradiol or a downstream metabolite is an endogenous ligand of PPAR γ . These results would also lead us to predict that estrogen would mimic thiazolidinediones in inhibiting aromatase expression in adipose stromal cells. Preliminary results have shown that estradiol in high concentrations does this (Fig. 3). Thus, there appears to be an important homeostatic mechanism operating in these cells to regulate the levels of estrogen biosynthesis in the context of adipocyte differentiation.

Testicular estrogen and spermatogenesis

Following our recent observations that male ArKO mice develop a progressive infertility such that by the age of 1 year most are infertile, we have examined the phenotype of these animals. We observed that the seminiferous tubules are grossly dysmorphic, with a complete absence of sperm and developmental arrest at the level of the round spermatids. In some cases multinucleate cells can be seen sloughing into the lumen. There is also marked Leydig cell hyperplasia, presumably as a consequence of the elevated

luteinizing hormone (LH) levels. In some cases, a normal tubule is seen side by side with a dysmorphic one. This phenotype is first noticed at around 20-23 weeks of age, and differs from that of the ERKO mice, which are infertile throughout life and have distended seminiferous tubules, apparently as a consequence of pressure build-up due to failure of fluid transport across the epithelium of the head of the epididymus. These observations point to important roles for intra-testicular estrogen production in male reproduction, and, in particular, spermatogenesis.

Clinical considerations

An important issue pertaining to the role of estrogen in the development of breast cancer in postmenopausal women is the relationship between HRT and breast cancer risk. A collaborative analysis of a large body of the available epidemiologic data which address this issue found that, despite the influence of estrogen produced locally on the development of breast cancer, systemic administration of estrogens plus progestins to postmenopausal women leads to at most a 1.35-fold increase in breast cancer risk (Collaborative Group on Hormonal Factors in Breast Cancer 1997). The reason for this small effect may have been revealed by the studies discussed here. Locally




Figure 3 Inhibition of aromatase activity of human adipose stromal cells by the PPAR ligands BRL 49653, and 15-deoxy- $\Delta^{12,14}$ -PGJ₂ (15d PGJ₂), in the presence of TNF α plus dexamethasone (Dex), or else oncostatin M (OSM) plus Dex. The inhibitory action of estradiol is also shown.

produced estrogen within the breast stimulates breast cancer development and is regulated by the mechanisms which have been discussed. The resulting intratumoral estradiol concentrations are one order of magnitude higher than in the circulating plasma (Pasqualini *et al.* 1996). Thus, the increase in circulating estrogen as a consequence of HRT may have little influence on intratumoral levels. The action of locally-produced estrogen is largely paracrine in nature, and mediated via the classical ER(s) or else via DNA adduct formation by quinone intermediates (Service 1998). Additionally, estrogens may have an autocrine or 'intracrine' action on adipose cells themselves whereby estradiol or a downstream metabolite generated within the cell site of synthesis may act by an alternative mechanism, possibly involving PPAR γ , to inhibit aromatase expression.

In conclusion, we believe that the results of recent studies reveal the significance of local estrogen production in the physiology and pathophysiology of elderly women and men, in particular its importance in the maintenance of bone mineralization and in the development of breast cancer, as well as its role in male reproduction. Local estrogen production may also play a role in the prevention of cardiovascular disease and in the maintenance of cognitive function. These studies not only throw light on the role of locally-produced estrogens in health and disease processes, but may also lead to new and hitherto unexpected modalities of therapy. This is already apparent from the observation that tumor-derived PGE₂ is the major factor stimulating local aromatase expression in the breast fat of cancer patients, which leads to the consideration that PG synthesis inhibitors such as aspirin and ibuprofen would be beneficial in breast cancer prevention or treatment (Zhao *et al.* 1996b), and indeed there is an epidemiologic study which supports this (Harris *et al.* 1996). Similarly, the observation that PPAR γ ligands, namely the thiazolidinediones, inhibit aromatase expression would suggest that these compounds, which are now available in the USA for the treatment of insulin-resistant diabetes, would also be beneficial in breast cancer prevention.

Acknowledgements

This work was supported by the Victorian Breast Cancer Research Consortium Inc., by Project Grant No. 981126 from the NH&MRC (Australia) and by Grant No. AGO8174 from the NIA (USA). The authors thank Faye Coates for skilled editorial assistance.

References

Agarwal VR, Bulun SE, Leitch M, Rohrich R & Simpson ER 1996 Use of alternative promoters to express the aromatase cytochrome P450 (*CYP19*) gene in breast adipose tissues of

cancer-free and breast cancer patients. *Journal of Clinical Endocrinology and Metabolism* **81** 3843-3849.

- Bayard F, Clamens S, Delsol G, Blaes N, Maret A & Faye JC 1995 Oestrogen biosynthesis, oestrogen metabolism and functional oestrogen receptors in bovine aortic endothelial cells. *Ciba Foundation Symposia* **191** 122-132.
- Bruch HR, Wolf L, Budde R, Romalo G & Schweikert HU 1992 Androstenedione metabolism in cultured human osteoblast-like cells. *Journal of Clinical Endocrinology and Metabolism* **75** 101-105.
- Bruning PF, Bonfrer JMG, van Noord PAH, Hart AAM, de Jong-Bakker M & Nooijen WJ 1992 Insulin resistance and breast-cancer risk. *International Journal of Cancer* **52** 511-516.
- Bulun SE & Simpson ER 1994 Competitive RT-PCR analysis indicates levels of aromatase cytochrome P450 transcripts in adipose tissue of buttocks, thighs, and abdomen of women increase with advancing age. *Journal of Clinical Endocrinology and Metabolism* **78** 428-432.
- Carani C, Qin K, Simoni M, Fanstini-Fustini M, Serpanti S, Boyd J, Korach K & Simpson ER 1997 Aromatase deficiency in the male: effect of testosterone and estradiol treatment. *New England Journal of Medicine* **337** 91-95.
- Collaborative group on hormonal factors in breast cancer 1997 Collaborative re-analysis of data from 51 epidemiological studies of 52705 women with breast cancer and 108441 women without breast cancer. *Lancet* **350** 1047-1059.
- Corbould AM, Judd SJ & Rodgers RJ 1998 Expression of types 1, 2 and 3. 17 Beta-hydroxysteroid dehydrogenase in subcutaneous abdominal and intra-abdominal adipose tissue of women. *Journal of Clinical Endocrinology and Metabolism* **83** 187-194.
- Davis SR, McCloud PI, Strauss BJG & Burger HG 1995 Testosterone enhances estradiol's effects on post-menopausal bone density and sexuality. *Maturitas* **21** 227-236.
- Fisher CR, Graves KH, Parlow AS & Simpson ER 1998 Characterization of mice deficient in aromatase (ArKO) due to targeted disruption of the *cyp19* gene. *Proceedings of the National Academy of Sciences of the USA* **95** 6965-6970.
- Harris RE, Nauboodim KK & Farrar WB 1996 Nonsteroidal anti-inflammatory drugs and breast cancer. *Epidemiology* **1** 203-205.
- Hemsell DL, Grodin JM, Brenner PF, Siiteri PK & MacDonald PC 1974 Plasma precursors of estrogen. II. Correlation of the extent of conversion of plasma androstenedione to estrone with age. *Journal of Clinical Endocrinology and Metabolism* **38** 476-479.
- Hotamisligil GS, Shargil NS & Spiegelman BM 1993 Adipose expression of tumor necrosis factor α : direct role in obesity-linked insulin resistance. *Science* **259** 87-91.
- Huang Z, Hankinson SE, Colditz GA, Stampfner MJ, Hunter DJ, Manson JE, Hennekens CH, Rosner B, Speizer FE & Willett WC 1997 Dual effects of weight and weight gain on breast cancer risk. *JAMA* **278** 1407-1411.
- Jakob F, Hormann D, Seufert J, Schneider D & Kohrle J 1995 Expression and regulation of aromatase cytochrome P450 in THP-1 human myeloid leukemia cells. *Molecular and Cellular Endocrinology* **110** 27-33.

- Labrie F, Belanger A, Cusan L & Candas B 1997a Physiological changes in dehydroepiandrosterone are not reflected by serum levels of active androgens and estrogens but of their metabolites: intracrinology. *Journal of Clinical Endocrinology and Metabolism* **82** 2403-2409.
- Labrie F, Belanger A, Cusan L, Gomez JL & Candas B 1997b Marked decline in serum concentrations of adrenal C19 sex steroid precursors and conjugated androgen metabolites during aging. *Journal of Clinical Endocrinology and Metabolism* **82** 2396-2402.
- Labrie F, Belanger A, Luu-The V, Labrie C, Simond J, Cusan L, Gomez JL & Candas B 1998 DHEA and the intracrine formation of androgens and estrogens in peripheral target tissues: its role during aging. *Steroids* **63** 322-328.
- Lubahn DB, Moyer JS, Golding TS, Couse JF, Korach KS & Smithies O 1993 Alteration of reproductive function but not prenatal sexual development after insertional disruption of the mouse estrogen receptor gene. *Proceedings of the National Academy of Sciences of the USA* **90** 11162-11166.
- Melton LJ 1997 Epidemiology of spinal osteoporosis. *Spine* **22** 2S-11S.
- Morishima A, Grumbach MM, Simpson ER, Fisher C & Qin K 1995 Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. *Journal of Clinical Endocrinology and Metabolism* **80** 3689-3698.
- Murakami H, Sasano H, Satoh A, Satomi S, Nagura H & Harada N 1998 Aromatase in human aortic tissue. *79th Annual Meeting of the Endocrine Society, Minneapolis, MN, USA* p212 (Abstract).
- Naftolin F, Ryan KJ, Davies IJ, Reddy VV, Flores F, Petro Z, Kuhn M, White RJ, Takaoka Y & Wolin L 1975 The formation of estrogens by central neuroendocrine tissues. *Recent Progress in Hormone Research* **31** 295-319.
- Nitta H, Bunick D, Hess RA, Janulis L, Newton SC, Millette CF, Osawa Y, Shizuta Y, Toda K & Bahr JM 1993 Germ cells of the mouse testis express P450 aromatase. *Endocrinology* **132** 1396-1401.
- Oz OK, Fisher CR, Graves KN, Nanu L, Zerwekh J & Simpson ER 1998 Alterations in bone histomorphometry and growth in aromatase deficient (ArKO) mice. *80th Annual Meeting of the Endocrine Society, New Orleans, LA, USA* p61 (Abstract).
- Pasqualini JR, Chetrite G, Blacker C, Feinstein MC, De la Londe L, Talbi M & Maloche C 1996 Concentrations of estrone, estradiol and estrone sulfate and evaluation of sulfatase and aromatase activities in pre- and postmenopausal breast cancer patients. *Journal of Clinical Endocrinology and Metabolism* **81** 1460-1464.
- Raisz LG, Wiita B & Artis A 1995 Comparison of the effects of estrogen alone and estrogen plus androgen on biochemical markers of bone formation and resorption in postmenopausal women. *Journal of Clinical Endocrinology and Metabolism* **81** 37-43.
- Sasano H, Frost AR, Saitoh R, Harada N, Poutanen M, Vihko R, Bulun SE, Silverberg SG, Nagura H 1996 Aromatase and 17 β -hydroxysteroid dehydrogenase Type 1 in human breast carcinoma. *Journal of Clinical Endocrinology and Metabolism* **81** 4042-4046.
- Sasano H, Uzuki M, Sawai T, Nagura H, Matsunaga G, Kashimoto O & Harada N 1997 Aromatase in human bone tissue. *Journal of Bone and Mineral Research* **12** 1416-1423.
- Service RF 1998 New role for estrogen in cancer? *Science* **279** 1631-1633.
- Shozu M & Simpson ER 1998 Aromatase expression of human osteoblast-like cells. *Molecular and Cellular Endocrinology* **139** 117-129.
- Shozu M, Zhao Y & Simpson ER 1997 Estrogen biosynthesis in THP1 cells is regulated by promoter switching of the aromatase (CYP19) gene. *Endocrinology* **138** 5125-5135.
- Siiteri PK & MacDonald PC 1973 Role of extraglandular estrogen in human endocrinology. In *Handbook of Physiology*, Vol 2, pp 619-629. Eds RO Greep & EB Astwood. Washington DC, USA: American Physiological Society.
- Simpson ER & Davis SR 1998 Why do the clinical sequelae of estrogen deficiency affect women more frequently than men? *Journal of Clinical Endocrinology and Metabolism* **83** 2214.
- Simpson ER, Zhao Y, Agarwal VR, Michael MD, Bulun SE, Hinshelwood MM, Graham-Lorence S, Sun T, Fisher CR & Mendelson CR 1997 Aromatase expression in health and disease. *Recent Progress in Hormone Research* **52** 185-213.
- Smith EP, Boyd J, Frank GR, Takahashi H, Cohen RM, Specker B, Williams TC, Lubahn DB & Korach KS 1994 Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. *New England Journal of Medicine* **331** 1056-1061.
- Spiegelman BM 1998 PPAR-gamma: adipogenic regulator and thiazolidinedione receptor. *Diabetes* **47** 507-514.
- Tsai-Morris CH, Aquilana DR & Dufau ML 1985 Cellular localization of rat testicular aromatase activity during development. *Endocrinology* **116** 38-46.
- Uysal KT, Wiesbrock SM, Marino MW & Hotamisligil GS 1997 Protection from obesity-induced insulin resistance in mice lacking TNF α function. *Nature* **389** 610-614.
- Yaffe K, Sawaya G, Lieberburg L & Grady D 1998 Estrogen therapy in postmenopausal women: effects on cognitive function and dementia. *JAMA* **279** 688-695.
- Zhao Y, Nichols JE, Bulun SE, Mendelson CR & Simpson ER 1995 Aromatase P450 gene expression in human adipose tissue: role of a Jak/STAT pathway in regulation of the adipose-specific promoter. *Journal of Biological Chemistry* **270** 16449-16457.
- Zhao Y, Nichols JE, Valdez R, Mendelson CR & Simpson ER 1996a Tumor necrosis factor- α stimulates aromatase gene expression in human adipose stromal cells through use of an activating protein-1 binding site upstream of promoter I.4. *Molecular Endocrinology* **10** 1350-1357.
- Zhao Y, Agarwal VR, Mendelson CR & Simpson ER 1996b Estrogen biosynthesis proximal to a breast tumor is stimulated by PGE₂ via cyclic AMP, leading to activation of promoter II of the CYP19 (aromatase gene). *Endocrinology* **150** S51-S57.