Hormone therapy in epithelial ovarian cancer

A P Makar

Departments of Gynecology and Oncology, The Middelheim Hospital, Lindendreef 1 2020 Antwerp, Belgium

Abstract

Although epidemiologic studies, animal experiments and receptor studies have shown that not only normal ovaries but also many malignant ovarian tumors can be considered as endocrine related and hormone dependent, the place of hormonal therapy in the management of patients with ovarian cancer remains unsettled. Most trials of hormonal treatment in ovarian cancer have been retrospective, involved only limited numbers of patients, and lacked important patient-related data and information pertaining to tumor characteristics. In addition, a variety of hormonal preparations with different degrees of potency and in different dosages were included in these studies. A literature review shows that response to hormonal therapy even in a preterminal setting, is modest, with about 8% objective response but almost no side effects. In a similar patient setting, more toxic therapeutic agents do not yield a better response. The place of hormonal therapy in the management of patients with epithelial ovarian cancer needs more thorough evaluation in well-designed randomized trials.

Endocrine-Related Cancer (2000) 7 85–93

Introduction

Ovarian cancer is the leading cause of death from all gynecologic malignancies. The disease affects elderly or middle aged women and the greatest incidence rates are reported in North America and Northern Europe, particularly in Scandinavia (Parkin 1989, Adami *et al.* 1990, Trope & Makar 1991, Pettersson 1994). The incidence rate in Norway was 19.3 per 100 000 in 1989. The incidence of ovarian cancer increases with age; it is relatively rare in women younger than 30 years, with only 1.5 new cancers/100 000 women per year in the 20–29 years age group, but beyond this age the incidence increases rapidly to a rate of 49/100 000 in women in the 60–69 years age group (Mant & Vessey 1995).

Ovarian cancer is characterized by its insidious onset and lack of early specific symptoms. About two-thirds of patients with ovarian cancer will present in International Federation of Gynecology and Obstetrics (FIGO) stages III and IV, having widespread tumor dissemination in the abdominal cavity, with or without varying degrees of pleural effusion. The prognosis of these patients remains poor, with 5-year actuarial survival of 23% and 14% for FIGO stages III and IV, respectively (Pettersson 1994, Makar *et al.* 1995). Management of these patients consists mainly of primary debulking surgery followed by platinum-based chemotherapy (Makar 1995, Makar *et al.* 1995).

Over the past several years, combinations of platinum and taxol became standard in the primary management of

advanced ovarian cancer (Caldas & McGuire 1993). Despite a primary response rate of up to 70%, the majority of patients suffer relapse (Neijt 1995). The probability of response to second-line chemotherapy depends mostly on the relapse-free interval. Patients with disease relapse during a regimen of platinum-based chemotherapy or within 6 months after its discontinuation are considered to be platinum resistant. Unfortunately, regimens used as second-line therapy in such situations have a low response rate and are associated with severe side effects that negatively influence the quality of life and necessitate long stays in hospital and intensive palliative care. It is therefore important to explore alternative treatment modalities that have cost–benefit ratios favorable in terms of toxicity. Hormonal therapy perhaps best fulfils these requirements.

Epidemiologic studies, animal experiments and receptor studies show that not only normal ovaries but also many malignant ovarian tumors can be considered to be endocrine related and hormone dependent.

Epidemiological studies

Several case–control studies have found a decrease in risk of the epithelial ovarian cancer associated with pregnancy, breast feeding and use of oral contraceptives (OCs). This prompted the hypothesis of incessant ovulation, which holds that factors that suppress ovulation may reduce the risk of developing ovarian cancer (Fathalla 1971, Wu *et al.* 1988,

Franceschi 1989). Gwin *et al.* (1989) showed, in a multicenter population-based, case-control study, that the estimated relative risks of epithelial ovarian cancer were 0.6 (95% confidence interval 0.5 to 0.8) for women who have ever been pregnant, 0.6 (95% confidence interval 0.5 to 0.8) for women who have ever breast fed, and 0.5 (95% confidence interval 0.5 to 0.7) for women who have ever used oral contraceptives. The risk of ovarian cancer seemed to decrease with increasing duration of use of OCs and the protective effect of OCs was even noticed in ex-users (at least 15 years). The findings of Van Leeuwen & Rookus (1989) were in agreement. The study by Wu *et al.* (1988) also showed that the risk of ovarian cancer tends to decrease with increasing age at menarche. Similarly, the risk of ovarian cancer was significantly decreased in women reporting life-long menstrual irregularities (Parazzini *et al.* 1989). Smith & Oi (1984) concluded that a group to be regarded at high risk should comprise women aged 45 years and older who are nulliparous, or whose first pregnancy occurred after the age of 30, who experienced menopause after age 55 years, or who had 40 years or more of 'ovulatory age'. However, the case-control study of Risch *et al.* (1983) was able only partly to confirm the theory of incessant or extravagant ovulation as a possible mechanism in the pathogenesis of ovarian cancer. These workers used a logistic regression method to test the hypothesis that equal periods of anovulation, regardless of cause, produced the same reduction in ovarian cancer risk. They found that the amounts of anovulatory time attributable to different exposures did not completely account for their protective effects. They suggested that the explanation of such a correlation cannot be based merely on hormonal suppression (Risch *et al.* 1983).

During the past decades, with new advances in fertility treatment, many questions have arisen regarding the effect of fertility drugs on the pathogenesis of ovarian cancer. The answer remains unclear. Clomiphene does not seem to be associated with any increased risk of ovarian cancer when used for fewer than 12 cycles. The relationship between gonadotropin use and ovarian cancer risk is less clear because of the small numbers of women exposed in the studies and short follow-up times. Prospective cohort studies are necessary, including more women exposed to clomiphene and gonadotropin and with correction for all the necessary confounding factors such as parity and with longer follow-up times. The putative risks must, however, be discussed with all patients before fertility treatment is commenced (Whittemore *et al.* 1992, Whittemore 1993, Mosgaard *et al.* 1997, Parazzini *et al.* 1997).

Experimental studies

The results of receptor and *in vitro* studies have also shown that ovarian cancer is an endocrine-related cancer (Amin *et*

al. 1987, Galtier-Dereure *et al.* 1992, Langdon *et al.* 1994a, b, Hua *et al.* 1995, Miller & Langdon 1997).

Slotman & Rao (1988), in a comprehensive analysis of 52 studies, reported the presence of estrogen (ER) and androgen receptors (AR) in the majority of primary ovarian cancers (63% and 69% respectively). Progesterone receptors (PgR) were also found in about 50% of tumors. Similarly, 88% of ovarian cancers were found to express glucocorticoid receptors (Galli *et al.* 1981). There has been inconsistency regarding the presence of greater levels of hormonal receptors, particularly PgR, in ovarian adenocarcinomas of the endometrioid type and in postmenopausal women (Sutton *et al.* 1986).

Binding sites for gonadotropic hormones and gonadotropic releasing hormone have also been detected in ovarian tumors. The receptors for luteinizing hormone-releasing hormone (LHRH) were found in nearly 80% of human ovarian cancers (Simons *et al.* 1983).

Estrogen-stimulated growth has been demonstrated in several cell lines characterized by an ER content greater than 23–30 fmol/mg protein (Nash *et al.* 1989b, Langdon *et al.* 1994b, Miller & Langdon 1997). Cell lines with lower ER concentrations appear unresponsive (Langdon *et al.* 1994b). As in ER-positive breast cancer models, 17- β estradiol (E2) regulates expression of the number of proteins associated with growth and invasion. On exposure to E2, the ER content of ER-positive cell lines is downregulated and the PgR receptor content is increased, both in culture and *in vivo* (Hamilton *et al.* 1984, Geisenger *et al.* 1989, Nash *et al.* 1989b, Miller & Langdon 1997). Expression of transforming growth factor (TGF)- α is also increased and, consistent with the action of this growth factor, there is eventual downregulation of the epidermal growth factor (EGF) receptor (Simpson *et al.* 1996, Miller & Langdon 1997). Other growth factor-mediated events include increased expression of TGF- β and modulation of several insulin-like growth factor (IGF) binding proteins (Nash *et al.* 1989a, Kyrwicki *et al.* 1993, Miller & Langdon 1997). Estrogen also activates the early growth response genes c-myc and c-fos (Hua *et al.* 1995, Miller & Langdon 1997). Mutations in the putative tumor suppressor gene BRCA1 are strongly associated with familial ovarian cancer and expression of this gene was found to be regulated by estrogen in an ovarian cancer cell line (Miller & Langdon 1997, Romagnolo *et al.* 1996). E2 also upregulates expression of procathepsin D, a protease likely to be involved in invasion and metastases. Overexpression of c-erbB-2 and cathepsin D in cancer cell lines is found to override estrogen control, leading to estrogen resistance (Hua *et al.* 1995, Miller & Langdon 1997).

Antiestrogens, including tamoxifen, have been found to inhibit estrogen-stimulated growth in ER-positive cell lines and xenografts. However, as in the case of endometrial cancer, tamoxifen also showed a growth-stimulating effect

under certain circumstances (Langdon *et al.* 1994a, Miller & Langdon 1997). Tamoxifen and ICI 182,780 (steroidal pure antiestrogen) also had a significant apoptotic effect even on ER-negative ovarian cancer cell lines, and this was found to have a dose-response relationship (Ercoli *et al.* 1998).

Kikuchi *et al.* (1993) showed that the presence of tamoxifen or clomiphene potentiated the antiproliferative effect of *cis*-diaminedichloroplatinum (CDDP) in ER-negative CDDP-resistant cancer cells by enhancing CDDP uptake in ovarian cancer cells by 30–80%. Similar results were obtained by McClay *et al.* (1994), who showed that tamoxifen delays the development of resistance to cisplatin in human melanoma and ovarian cancer cell lines.

Animal experiments have also shown LHRH agonists to have an antiproliferative effect on ovarian cancer cells (Peterson & Zimniski 1990). Peterson *et al.* (1994) showed this response to be latent and transient. A study by Yano *et al.* (1994) showed that chronic exposure to LHRH antagonists reduced the concentration of receptors for EGF and IGF-I in tumor cell membrane. This phenomenon is related to tumor growth inhibition, suggesting a direct antitumoral effect, rather than exclusively one of inhibition of the pituitary–gonadal axis.

A study by Miyazaki *et al.* (1997) compared the antiproliferative effect of LHRH (AN-152) with that of doxorubicin, in nude mice bearing LHRH-receptor-positive and -receptor-negative ovarian cancer cell lines. They showed a significant antiproliferative effect on LHRH-positive ovarian tumors in nude mice even with the lowest tested dose of AN-152. An equivalent dose of doxorubicin caused substantial mortality.

Although the data regarding effects of androgens and progestins are less extensive, Rose & Barnea (1996) reported the antiprogestin mifepristone to downregulate progesterone expression on human ovarian cancer cell lines. Mifepristone blocked cells in a G₀/G₁ phase and thus reduced the number of cells in the S phase. The efficacy of mifepristone was compared with that of taxol and tamoxifen in the same ovarian cancer cell line. Continuous exposure to tamoxifen resulted in a varied cytostatic response and in a transient change in the cell cycle. Taxol inhibited growth of some but not all cancer cell lines.

The data regarding glucocorticosteroids are controversial. Both dexamethasone and cortisol increase expression of c-erbB-2 by a post-transcriptional mechanism and inhibit production of the tumor marker CA 125 (Karlán 1988, Hua *et al.* 1995, Miller & Langdon 1997).

The postmenopausal ovary and steroid hormone production

The cortex of the postmenopausal ovary is thin. The remaining follicles are few and atretic. Granulosa cells have disappeared and are replaced by fibrocytes, and theca cells

seem to differentiate towards interstitial cells of the stroma. These cells are still capable of hypertrophy and of steroid production in response to LH or human chorionic gonadotrophin. In contrast to the premenopausal situation, no cyclic variation in steroid hormone production by the ovary occurs after the menopause (Greenblatt *et al.* 1976).

Although production of progesterone ceases after menopause, lower levels of progesterone were observed in oophorectomized women than in postmenopausal controls (Bakstrom *et al.* 1983). The postmenopausal concentrations of androstenedione are roughly comparable to those in the early follicular phase (smallest premenopausal plasma values) (Abraham *et al.* 1969, Mahlck 1989). Concentrations are also lowest in oophorectomized women.

Testosterone concentrations do not decrease after the menopause, possibly because of an increase in postmenopausal gonadotropin. Another reason could be the unchanged binding capacity of sex hormone binding globulin after menopause, as metabolism acts only on the unbound fraction (Judd *et al.* 1974a,b, Mahlck 1989).

Ovarian estrogen production ceases after the menopause, but some aromatase activity remains. The main origin of the postmenopausal circulating estrogen, essentially estrone, is found in the extraovarian aromatization of androstenedione in the skin and adipose tissue (Grodin *et al.* 1973).

Mahlck (1989) showed that, in postmenopausal patients with epithelial ovarian cancer, plasma concentrations of progesterone, androstenedione, estradiol and testosterone were directly correlated with tumor volume and FIGO stage. Positivity for PgRs and the preoperative plasma concentration of progesterone were found to be independent prognostic factors for survival (Mahlck 1989, Sevelde *et al.* 1990, Harding *et al.* 1990). As in the case of CA 125 serum measurements (Makar 1995), changes in hormonal concentrations (with the exception of testosterone) reflected the response of ovarian cancer to therapy. The combined measurement of progesterone and androstenedione also allowed detection of disease relapse earlier than did clinical means, in the majority of patients (Mahlck 1989).

Contrary to these findings, in other studies no consistent correlation between the presence of steroid receptors and response to hormonal therapy was observed (From *et al.* 1991, Slotman & Rao 1988). However, more sensitive and specific assays of hormone receptors are needed, as defective processing mechanisms and heterogeneity of the receptors within the tumor may influence the results (O'Brien *et al.* 1981). Burger *et al.* (1999) showed a weak correlation between immunohistochemical and biochemical analyses of hormone receptors in ovarian tumors. This discrepancy was explained by heterogeneity of the receptor within the tumor and the presence of progesterone and androgen in the stromal section of the tumor. The study also showed that ERs are mostly present in the epithelial component (Burger *et al.* 1999).

Patient trials

Hormonal therapy for epithelial ovarian cancer has resulted in uneven but consistent responses. Unfortunately, most trials of hormonal treatment of ovarian cancer have included only limited numbers of patients and were retrospective. In addition, important patient characteristics, including histologic type and tumor grade, type and extent in addition to the response to previous treatments, were not critically defined in the majority of cases. Furthermore, most reports implied the use of hormonal therapy in a setting in which almost every line of standard chemotherapy had failed. This last observation is important, as the clinical efficiency of salvage chemotherapy in ovarian cancer is strongly influenced by the presence of platinum resistance, indexed as previous response to platinum-containing chemotherapy and a treatment-free interval between the last cytotoxic drug and the initiation of second line chemotherapy. It is therefore essential in considering any salvage program to know how effective this treatment is in patients considered clinically to be platinum refractory.

The earliest suggestion that ovarian cancer might be hormone sensitive was from a study published by Long & Evans in 1963. Using diethylstilbestrol in 14 patients with advanced ovarian cancer, four (28%) achieved a partial response. Estrogen might also decrease serum gonadotropin concentrations by a mechanism of negative feedback inhibition. This is in agreement with findings of epidemiologic studies suggesting a reduction in the death rate from ovarian cancer among women younger than 55 years to be consistent with a protective effect of the oral contraceptive pill (Mant & Vessey 1995). Estrogen replacement therapy was equally found to have no adverse effects in patients with ovarian cancer (Guidozzi & Paponte 1999).

Progestational agents have been the most extensively studied, although there has been inconsistency regarding doses and routes of administration (Bergquist *et al.* 1981, Rendina *et al.* 1982, Geisler 1983, 1985, Trope *et al.* 1982, Thigpen *et al.* 1984, Van Der Vange *et al.* 1995). The response rate in a series with adequate numbers of patients and response criteria was about 7% (Van Der Vange *et al.* 1995). No dose-response relationship was consistently found and the trial by Geisler (1983), using high doses of megestrol acetate (800 mg/day for 30 days, then 400 mg/day) and showing an objective response of 43%, could not be confirmed by others (Sikic *et al.* 1986, Ahlgren *et al.* 1993, Wilailak *et al.* 1999). Although most of the reported series found no correlation between response and histologic types or grades, some overviews suggest more response in cases of serous and endometrioid adenocarcinomas of the ovary when given as first-line agents or after only a single alkylating agent (Bergquist *et al.* 1981, Rendina *et al.* 1982, Thigpen *et al.* 1984, Geisler 1985). Estrogen has the ability to induce PgRs, and was used in two different doses to

enhance treatment by progestational agents (Jolles *et al.* 1983). The results showed no difference between the two dosages, and the overall response rate was 14% (9/65). A similar regimen was used by From *et al.* (1991), who observed a partial response in 17% of their patients.

Perhaps the first report of clinical activity of antiestrogens in ovarian cancer was that by Myers *et al.* (1981). Tamoxifen has been the most used antiestrogen: the literature contains reports of more than 500 patients with advanced ovarian cancer who were treated with this drug (Schwartz *et al.* 1982, 1989, Shirley *et al.* 1985, Slevin *et al.* 1986, Beecham *et al.* 1987, Weiner *et al.* 1987, Markman *et al.* 1996, Marth *et al.* 1997). The reported rates of response to tamoxifen varied between 0 and 28% and the overall response rate was of the order of 8%. However, a significant percentage of patients (20–75%) benefitted from disease stabilization for some time. As in the case of trials with progestational therapy, most of the patients in these trials also were at a 'pre' terminal stage of disease and were refractory to different lines of treatment.

Markman, in 1996, updated a Gynecologic Oncology Group study evaluating tamoxifen in patients with advanced ovarian cancer after stratifying patients according to whether or not they were refractory to platinum. In 77 patients in the platinum-refractory group, a clinical response rate of 13% was described. The response rate in patients who were platinum sensitive did not differ markedly (15%). Stable disease was also found in more than 35% of patients with a median disease duration of 3 months. Van Der Velden *et al.* (1995) obtained a mean duration of stable disease of 11.5 months (4–30 months) when tamoxifen was started on the basis of increasing CA 125 serum concentrations without a clinically or radiologically detectable mass.

Although a Norwegian report (Marth *et al.* 1995) suggested a better response in endometrioid adenocarcinomas of the ovary, literature review reveals no apparent difference in histologic subtype, grade of tumor or hormone receptor values between responders and non-responders, and no correlation between response rate and tumor ER and PgR content was evident (Schwartz *et al.* 1982, 1989, Shirley *et al.* 1985, Slevin *et al.* 1986, Beecham *et al.* 1987, Weiner *et al.* 1987). Endometrioid tumors are known to have the best prognosis among all ovarian adenocarcinomas (Makar 1995, Makar *et al.* 1995). It remains an open question whether the better response observed with these tumors represents a therapeutic gain or merely reflects the behavior of biologically less aggressive tumors.

Only a few trials showed better responses with high doses of tamoxifen up to 80 mg (Ahlgren *et al.* 1993), and there is no convincing evidence that sequential use of tamoxifen and progestational agents improves response rates (Belinson *et al.* 1987, Jakobson *et al.* 1987).

The place of LHRH agonists in the management of ovarian cancer has also been evaluated, and the response rate observed in cases of chemotherapy-refractory ovarian cancer seems to be modest (Adelson & Reece 1993, Parmar *et al.* 1988, Bruckner & Motwani 1989, Kavanagh *et al.* 1989). Among the problems in evaluating the efficacy of such treatment is the enrolment of patients with expected poor outcome and the use of several different gonadotropin-releasing hormone (GnRH) analogs. It is unclear whether the various GnRH analogs possess similar antitumor potency *in vivo*, as marked differences in potency were observed in *in vitro* studies (Adelson & Reece 1993). In a review of the literature, Adelson & Reece (1993) reported statistically significant differences in the number of responders among patients treated with leuprolide as compared with those receiving goserelin.

In two reports including 59 patients, an objective response rate of 15% was observed in patients with advanced-stage disease (Parmar *et al.* 1988, Kavanagh *et al.* 1989); stable disease was reported in 12%. The study by Parmar *et al.* (1988), showed a mean duration of response of 10 months, and the median duration of response reported by Kavanagh's group was 52 weeks. The findings of Kavanagh *et al.* (1989) also suggested a greater response rate among patients with highly differentiated tumors. Van Der Vange *et al.* (1995), reported a complete response using LHRH agonists for 38 months in patients with highly differentiated ovarian tumors. Furthermore, a prospective analysis reported 12% partial responses (clinical or radiological) in 32 patients with advanced refractory ovarian cancer who were treated with leuprolide acetate (Marinaccio *et al.* 1996). The mean duration of response was 8.7 weeks and all responders showed a significant decrease in CA 125 concentrations. In addition, 15% of patients had stable disease for a mean time of 5.2 months. A significant difference in survival was observed between responders and non-responders ($P < 0.05$). All patients included in this study had either moderately or poorly differentiated tumors.

At least two studies with a total of 87 patients with advanced ovarian cancer, however, revealed no objective response to LHRH analogs (Scambia *et al.* 1990, Jager *et al.* 1995). The majority of patients in the report by Scambia's group had poorly differentiated tumors; the paper by Jager *et al.* (1995) did not report tumor grade. Furthermore, the addition of GnRH analogs to first-line platinum-based chemotherapy did not seem to improve the outcome of patients with advanced ovarian cancer in the prospective randomized double-blind study including 135 patients with FIGO stages III and IV published by Emons *et al.* (1996). Similar results were reported by Erickson *et al.* (1994) and Falkson *et al.* (1996).

Alternative endocrine approaches in the hormonal treatment of ovarian cancer include the use of androgens, antiandrogens, and aminoglutethimide. Androgens have been

used unsuccessfully (Kavanagh *et al.* 1987); negative results were also reported with aminoglutethimide (Rothschild *et al.* 1987, Ahlgren *et al.* 1993). From the theoretical point of view, antiandrogen therapy seems to be helpful. Decreased activity of aromatase enzyme in ovarian tumors leads to a local accumulation of androgens, as they are not converted to estrogen. As ovarian carcinomas without aromatase activity are AR-positive, androgen may influence the tumor cells through this receptor (Van Der Vange *et al.* 1995). This said, the literature contains few results suggesting beneficial effects with antiandrogen therapy (Slotman & Rao 1988, Van Der Vange *et al.* 1995).

Summary

Ovarian cancer is an endocrine-related cancer as shown by epidemiological, receptor and *in vitro* studies. In postmenopausal patients with epithelial ovarian cancer, the concentration of plasma progesterone has an independent prognostic significance and correlates with tumor volume, FIGO stage and response to chemotherapy. Furthermore, combined determination of progesterone and androstenedione concentrations allows early detection of disease relapse.

In contrast to findings in breast and endometrial cancers, receptor status in ovarian cancer was not predictive of hormonal response. Wide variations in the level of receptors in different cells from the same ovarian cancer have been observed, and the level of receptors in metastases appears to be lower than that in the primary tumor. The average magnitude of receptor concentration in ovarian cancer cells is lower than in breast or endometrial cancer cells, and ER concentrations of more than 100 fmol/mg – which are relatively common in breast cancer and have the greatest probability of response – are relatively uncommon in ovarian cancer (Miller & Langdon 1997). Although experimental studies have shown manifest antiproliferative activity in response to hormonal therapy that is comparable to that achieved with platinum, anthracycline or taxol, even in the presence of ER-negative cell lines, response rates in patient trials remain modest. Also, the addition of hormonal therapy to standard first-line chemotherapy failed to improve either response or progression-free survival in a limited number of trials. However, it is unclear whether the marginal efficiency of endocrine therapies (at best around 10% of objective responses in phase II trials) can be substantiated in controlled trials. For ethical reasons controlled trials that evaluate the value of single agent hormonal therapy as first-line treatment are difficult to perform in ovarian carcinoma.

A large number of trials have evaluated the place of hormonal therapy in cases of disease relapse (mostly in patients at a preterminal stage of disease). The inconsistency in defining patients and tumor characteristics included in these trials, the wide variations in the dosages of drugs used

(particularly in case of progestational agents) and the variety of preparations used that had different potency (particularly in case of GnRH analogues), are all factors that explain why the correct place of hormonal therapy in the management of ovarian cancer remains poorly evaluated, notwithstanding an objective response of around 8% and stable disease in a significant number of patients. In addition to the critical need to identify new drugs that are active in combatting ovarian cancer, it is also important to develop rational therapeutic strategies for the relapsed/refractory setting that focus on optimizing quality of life. The 8% objective response rate of tamoxifen in platinum-refractory ovarian cancer establishes this agent as equaling the demonstrated antitumor effectiveness of more toxic regimens such as hexamethylmelamine and ifosfamide. In view of the favorable toxicity profile of tamoxifen, certainly as compared with alternative cytotoxic chemotherapeutic drugs, it is not difficult to justify treating an individual with advanced ovarian cancer with this agent despite being unable to 'prove' that the 'stable disease state' is the direct result of treatment with this antihormone agent.

Finally, future studies are needed to evaluate the place of tamoxifen or other hormonal therapy in different settings in ovarian cancer such as:

1. Consolidation therapy for patients with pathological complete response after first-line chemotherapy. Hormonal therapy can be randomized with placebo.
2. Therapy for asymptomatic patients with an increasing CA 125 antigen concentration after first- or second-line chemotherapy.
3. Activity in patients previously treated with paclitaxel.
4. Identification of subsets of patients with less aggressive tumors who might derive maximum benefit from hormonal therapy, such as those with highly differentiated tumors, diploid tumors, tumors with c-erbB-2 expression, or tumors with high steroidal receptor content (Iversen 1988, Berchuck *et al.* 1990, Kaern *et al.* 1990, Makar *et al.* 1994, Van Der Vange *et al.* 1995, Miller & Langdon 1997).

References

- Abraham GE, Lobotsky J & Lloyd CW 1969 Metabolism of testosterone and androstenedione in normal and ovariectomized women. *Journal of Clinical Investigation* **48** 696–703.
- Adami AO, Bergstrom R, Persson I & Sparen P 1990 The incidence of ovarian cancer in Sweden 1960–1984. *American Journal of Epidemiology* **132** 446–452.
- Adelson MD & Reece MT 1993 Effects of gonadotropin-releasing hormone analogues on ovarian epithelial tumors. *Clinical Obstetrics and Gynecology* **36** 690–700.
- Ahlgren JD, Ellison NM, Gottlieb RJ, Laluna F, Lokich JJ, Sinclair PR, Ueno W, Wampler GL, Yeung KY, Alt D & Fryer JG 1993 Hormonal palliation of chemoresistant ovarian cancer: three consecutive phase III trials of the mid-atlantic oncology program. *Journal of Clinical Oncology* **11** 1957–1968.
- Amin W, Karlan BY & Littlefield BA 1987 Glucocorticoid sensitivity of OVCA 433 human ovarian carcinoma cells: inhibition of plasminogen activators, cell growth and morphological alterations. *Cancer Research* **47** 6040–6045.
- Bakstrom T, Mahlck CG & Kjellgren O 1983 Progesterone as a possible tumor marker for 'non-endocrine' ovarian malignant tumors. *Obstetrics and Gynecology* **16** 129–138.
- Beecham J, Blessing J, Creasman W & Hatch K 1987 The role of tamoxifen as second line therapy in advanced ovarian cancers evaluated for receptor status and tumor grade. *Proceedings of the International Gynecological Cancer Society* **1** 117.
- Belinson JL, McClure M & Badger G 1987 Randomized trial of megestrol acetate vs megestrol acetate/tamoxifen for the management of progressive or recurrent epithelial ovarian carcinoma. *Gynecology and Oncology* **28** 151–155.
- Berchuck A, Kamel A, Whitaker R, Kerns B, Olt G, Kinne GR, Soper JT, Dodge R, Clark Pearson & Merks P 1990 Overexpression of HER-2/neu is associated with poor survival in advanced epithelial ovarian cancer. *Cancer Research* **50** 4087–4091.
- Bergquist A, Kullander S & Thorell J 1981 A study of oestrogen and progesterone cytosol receptor concentration in benign and malignant ovarian tumors treated with medroxy progesterone acetate. *Acta Obstetrica et Gynaecologica Scandinavica* **101** 75–81.
- Bruckner HW & Motwani BT 1989 Treatment of advanced refractory ovarian carcinoma with a gonadotropin releasing hormone analogue. *American Journal of Obstetrics and Gynecology* **161** 1216–1218.
- Burger CW, van Doorn HC, van der Valk P & Bonfrer HMG 1999 Oestrogen, progesterone and androgen receptors in ovarian neoplasm; correlation between immunohistochemical and biochemical receptor analysis. *International Journal of Gynecological Cancer* **9** 17.
- Caldas C & McGuire WP III 1995 Paclitaxel (taxol) therapy in ovarian carcinoma. *Seminars in Oncology* **20** 50–56.
- Emons G, Ortmann O, Teichert HM, Fassl H, Lohrs U, Kullander S, Kauppila A, Ayalon D, Schally A & Oberheuser F 1996 Luteinizing hormone-releasing hormone agonist triptorelin in combination with cytotoxic chemotherapy in patients with advanced ovarian carcinoma. *Cancer* **78** 1452–1460.
- Ercoli A, Scambia G, Fattorossi A, Paspaglio G, Battaglia A, Cicchillitti L, Malorni W, Rainaldi G, Benedetti P & Mancuso S 1998 Comparative study of the induction of cytostasis and apoptosis by ICI 182,780 and tamoxifen in an estrogen receptor negative ovarian cancer cell line. *International Journal of Cancer* **76** 47–54.
- Erickson LD, Hartmann LC, Su JQ, Nielsen SNJ, Pfeifel DM, Goldberg RM, Levi R & Stanope CR 1994 Cyclophosphamide, cisplatin, and leuprolide acetate in patients with debulked stage III or IV ovarian carcinoma. *Gynecology and Oncology* **54** 196–200.
- Falkson CI, Falkson HC & Falkson G 1996 Cisplatin versus cisplatin plus d-Trp-6LHRH in the treatment of ovarian cancer: a pilot trial to investigate the effect of the addition of GnRH analogue to cisplatin. *Oncology* **53** 313–317.
- Fathalla MF 1971 Incessant ovulation – a factor in ovarian neoplasia? *Lancet* **ii** 163.
- Franceschi S 1989 Reproductive factors and cancers of the breast, ovary and endometrium. *European Journal of Cancer and Clinical Oncology* **25** 1933–1943.

- From GL, Freedman RS, Fritsche HA, Atkinson EN & Scott W 1991 Sequentially administered ethinyl and medroxyprogesterone acetate in the treatment of refractory epithelial ovarian carcinoma in patients with positive estrogen receptors. *Cancer* **68** 1885–1889.
- Galli MC, De Giovanni C & Nicoletti G, Grilli S, Nanni P, Prodi G *et al.* 1981 The occurrence of multiple steroid hormone receptors in disease free and neoplastic human ovary. *Cancer* **47** 1297–1302.
- Galtier-Dereure F, Capony F, Maudelonde T & Rochefort H 1992 Estradiol stimulates cell growth and secretion of precathepsin D and a 120-kilodalton protein in the human ovarian cancer cell line BG-1. *Journal of Clinical Endocrinology and Metabolism* **75** 1497–1502.
- Geisenger KR, Kutte TE, Pettenati MJ, Welander CE, Dennard Y, Collins LA & Berens ME 1989 Characterization of a human ovarian carcinoma cell line with estrogen and progesterone receptors. *Cancer* **63** 280–288.
- Geisler H 1983 Megestrol acetate for the palliation of advanced ovarian carcinoma. *Obstetrics and Gynecology* **61** 95–98.
- Geisler H 1985 The use of high-dose megestrol acetate in the treatment of ovarian cancer. *Seminars in Oncology* **11** 20.
- Greenblatt RB, Colle ML & Mahesh VB 1976 Ovarian and adrenal steroid production in the postmenopausal woman. *Obstetrics and Gynecology* **47** 383–387.
- Grodin JM, Siiteri PK & MacDonald PC 1973 Source of estrogen production in postmenopausal women. *Journal of Clinical Endocrinology and Metabolism* **36** 207–214.
- Guidozzi F & Paponte A 1999 Estrogen replacement therapy in ovarian cancer survivors. *International Journal of Gynecological Cancer* **9** 4.
- Gwin ML, Lee N, Rhodes PH, Layde PM & Rubin G 1989 Pregnancy, breast feeding, and oral contraceptives and the risk of epithelial ovarian cancer. *Journal of Clinical Epidemiology* **43** 559–568.
- Hamilton TC, Behrens BC, Louie KG & Ozols RF 1984 Induction of progesterone receptor in human ovarian cancer. *Journal of Clinical Endocrinology and Metabolism* **59** 561–563.
- Harding M, Cowan S, Hole D, Cassidy L, Kitchener H, Davis J & Leake R 1990 Estrogen and progesterone receptors in ovarian cancer. *Cancer* **65** 486–491.
- Hua W, Christianson T, Roougoet C, Rochefort H & Clinton G 1995 SKOV3 ovarian carcinoma cells have functional estrogen receptors but are growth resistant to estrogen and antiestrogens. *Journal of Steroid Biochemistry and Molecular Biology* **55** 279–289.
- Iversen OE 1988 Prognostic value of flow cytometric DNA index .. in human ovarian cancer. *Cancer* **61** 971–975.
- Jager W, Sauerhrei W, Beck E, Maasen V, Stumpfe M, Meier W, Kunh W & Janicke F 1995 A randomized comparison of triptorelin and tamoxifen as treatment of progressive ovarian cancer. *Anticancer Research* **15** 2639–2642.
- Jakobsen A, Bertelsen K & Sell A 1987 Cyclic hormonal treatment in ovarian cancer. A phase II trial. *European Journal of Cancer and Clinical Oncology* **23** 915–916.
- Jolles CJ, Freedman RS & Jones LA 1983 Estrogen and progesterone therapy in advanced ovarian cancer: preliminary report. *Gynecology and Oncology* **16** 352–359.
- Judd HL, Judd GE, Lucas WE & Yen SSC 1974a Endocrine function of the postmenopausal ovary: concentration of androgens and estrogens in ovarian and peripheral vein blood. *Journal of Clinical Endocrinology and Metabolism* **39** 1020–1024.
- Judd HL, Lucas WE & Yen SSC 1974b Effect of oophorectomy on circulating testosterone and androstenedione levels in patients with endometrial cancer. *American Journal of Obstetrics and Gynecology* **118** 793–798.
- Kaern J, Trope CG, Kjørstad KE, Abeler V & Pettersen EO 1990 Cellular DNA contents as a new prognostic tool in patients with borderline tumors of the ovary. *Gynecology and Oncology* **38** 452–457.
- Karlan BY, Amin W, Casper SE & Littlefield BA 1988 Hormonal regulation of CA 125 tumor marker expression in human ovarian carcinoma cells: inhibition by glucocorticoids. *Cancer Research* **48** 3502–3506.
- Kavanagh JJ, Wharton JT & Roberts WS 1987 Androgen therapy in the treatment of refractory epithelial ovarian cancer. *Cancer Treatment Report* **71** 537–538.
- Kavanagh JJ, Roberts W, Townsend P & Hewitt S 1989 Leuprolide acetate in the treatment of refractory or persistent epithelial ovarian cancer. *Journal of Clinical Oncology* **7** 115–118.
- Kikuchi Y, Hirata J, Kita T, Imaizumi E, Tode T & Nagata I 1993 Enhancement of antiproliferative effect of cis-diaminedichloroplatinum (II) by clomiphene and tamoxifen in human ovarian cancer cells. *Gynecology and Oncology* **49** 365–372.
- Kyrwicki RF, Figueroa JA, Jacson JC, Wozelsky TW, Shimasaki S, von Hoff DD & Yee D 1993 Regulation of insulin-like growth factor binding proteins in ovarian cancer cells by oestrogen. *European Journal of Cancer* **29A** 2015–2019.
- Langdon SP, Crew AJ, Ritchie AA, Muir M, Wakeling A, Smyth JF & Miller WR 1994a Growth inhibition of oestrogen receptor-positive human ovarian carcinoma by anti-oestrogens *in vitro* and in a xenograft model. *European Journal of Cancer* **30A** 682–686.
- Langdon SP, Hirst GL, Miller EP, Hawkins RA, Tesdale AL, Smyth JF & Miller WR 1994b The regulation of growth and protein expression by estrogen *in vitro*: a study of 8 human ovarian carcinoma cell lines. *Journal of Steroid Biochemistry and Molecular Biology* **50** 131–135.
- Long RTL & Evans AM 1963 Diethylstilbestrol as a chemotherapeutic agent for ovarian cancer. *Modern Medicine* **60** 1125–1127.
- McClay EF, Albright KD, Jones JA, Christen RD & Howell SB 1994 Tamoxifen delays the development of resistance to cisplatin in human melanoma and ovarian cancer cell lines. *British Journal of Cancer* **70** 449–452.
- Mahlck CG 1986 Plasma steroid hormones in women with epithelial ovarian carcinoma. Doctoral thesis. *Acta Obstetrica et Gynecologica Scandinavica (Suppl)* **137** 1–32.
- Makar AP 1995 Prognostic studies in cancer of the ovary and fallopian tube with emphasis on the CA 125 and c-erbB-2 oncogene. Summary of doctoral thesis. *Acta Obstetrica et Gynecologica Scandinavica* **74** 238–240.
- Makar AP, Kristensen GB, Nesland J, Borner OP & Trope CG 1994 The expression of c-erbB-2 (HER-2/neu) oncogene in patients with invasive ovarian malignancies. *International Journal of Gynecological Cancer* **4** 194–199.
- Makar AP, Baeklandt M, Trope CG & Kristensen GB 1995 The prognostic significance of residual disease, FIGO substage, tumor histology and grade in patients with FIGO stage III ovarian cancer. *Gynecology and Oncology* **56** 175–180.

- Mant JWF & Vessey MP 1995 Mortality trends in ovarian cancer. In *Epithelial Cancer of the Ovary*, edn 1, pp 1–12. Eds FG Lawton, JP Neijt & KD Swenerton. London: BMJ Publishing Group.
- Marinaccio M, D'Addario VD, Serrati A, Pinto V & Gagnazzo G 1996 Leuprolide acetate as a salvage therapy in relapsed epithelial ovarian cancer. *European Journal of Clinical Oncology* **17** 286–288.
- Markman M, Iseminger KA, Hatch KD, Creasman WT, Barnes W & Dubeshter B 1996 Tamoxifen in platinum refractory ovarian cancer. A Gynecologic Oncology Group ancillary report. *Gynecology and Oncology* **62** 4–6.
- Marth C, Sorheim N, Kaern J & Trope C 1997 Tamoxifen in the treatment of recurrent ovarian carcinoma. *International Journal of Gynecological Cancer* **7** 256–261.
- Miller WR & Langdon SP 1997 Steroid hormones and cancer: (II) lessons from experimental system. *European Journal of Surgical Oncology* **23** 72–83.
- Miyazaki M, Nagy A, Schally AV, Lamharzi N, Halmos G, Szepeshazi K, Groot K & Armatis P 1997 Growth inhibition of human ovarian cancers by cytotoxic analogues of luteinizing hormone-releasing hormone. *Journal of the National Cancer Institute* **89** 1803–1809.
- Mosgaard BJ, Lidegaard O & Andersen AN 1997 The impact of parity, infertility and the treatment with fertility drugs on the risk of ovarian cancer. *Acta Obstetrica et Gynaecologica Scandinavica* **76** 89–95.
- Myers AM, Moore GE & Major FJ 1981 Advanced ovarian carcinoma: response to antiestrogen therapy. *Cancer* **48** 2368–2370.
- Nash J, Hall L, Ozols R, Young R, Smyth J & Hamilton T 1989a Estrogenic regulation and growth factor expression in human ovarian cancer *in vitro*. *Proceedings of the American Association for Cancer Research* **30** 1189.
- Nash JD, Ozols RF, Smyth JF & Hamilton TC 1989b Estrogen and antiestrogen effects on the growth of human epithelial ovarian cancer *in vitro*. *Obstetrics and Gynecology* **73** 1009–1016.
- Neijt JP 1995 Optimum chemotherapy regimens, dose intensity, and new drugs. In *Epithelial Cancer of the Ovary*, edn 1, pp 186–209. Eds FG Lawton, JP Neijt & KD Swenerton. London: BMJ Publishing Group.
- O'Brien TJ, Hernandez W, Jernstrom PJ, Seymour BD, Morrow CP & Sykes JA 1981 Modulation of protein expression in endometrial carcinoma cells by *in vitro* exposure to estradiol and progesterone. *American Journal of Obstetrics and Gynecology* **139** 67–72.
- Parazzini F, La Vecchia C, Negri E & Gentile A 1989 Menstrual factors and the risk of epithelial ovarian cancer. *Journal of Clinical Epidemiology* **5** 443–448.
- Parazzini F, Negri E, La Vecchia C, Moroni S, Franceschi S & Crosignani PG 1997 Treatment of infertility and risk of invasive epithelial ovarian cancer. *Human Reproduction* **12** 2159–2161.
- Parkin MD 1989 Cancers of the breast, endometrium, and ovary: geographic correlations. *European Journal of Cancer and Clinical Oncology* **25** 1917–1925.
- Parmar H, Philips RH, Rustin G, Lightman SL & Shally AV 1988 Therapy of advanced ovarian cancer with d-Trp-6-LH-RH (decaptyl) microcapsules. *Biomedicine and Pharmacotherapy* **42** 531–538.
- Peterson CM & Zimmiski SJ 1990 A long-acting gonadotropin releasing hormone agonist inhibits the growth of a human ovarian epithelial carcinoma (BG-1) heterotransplanted in the nude mouse. *Obstetrics and Gynecology* **76** 264–267.
- Peterson CM, Jolles CJ, Carrell DT, Straight RC, Jones KP, Poulson AM Jr & Hatasaka H 1994 GnRH agonist therapy in human ovarian epithelial carcinoma (OVCAR-3) heterotransplanted in the nude mouse is characterized by latency and transience. *Gynecology and Oncology* **52** 26–30.
- Pettersson F 1994 *Annual Report on the Result of Treatment in Gynecological Cancer*. Stockholm, Sweden: FIGO **22** 83–102.
- Rendina GM, Donadio C & Giovannini M 1982 Steroid receptors and progestin therapy in ovarian endometrioid carcinoma. *European Journal of Gynaecological Oncology* **3** 241–246.
- Risch HA, Weiss NH & Lyon JL 1983 Events of reproductive life and the incidence of epithelial ovarian cancer. *American Journal of Epidemiology* **117** 1128–1139.
- Romagnolo D, Annab LA, Risinger JL, Terry LA, Lyon TT & Barrett JC 1996 Regulation of expression of BRCA-1 cancer cells. *Proceedings of the American Association for Cancer Research* **37** 516.
- Rose FV & Barnea ER 1996 Response of human ovarian carcinoma cell lines to antiprogestin mifepristone. *Oncogene* **12** 999–1003.
- Rothschild N, Gray R, Smith L, Matelski H & Ahlgren J 1987 Phase II study aminoglutethimide in advanced ovarian cancer. *Proceedings of the American Society of Clinical Oncology* **6** 115.
- Scambia G, Benedetti Panici P, Greggi S, Baiocchi G, Perrone L & Mancuso S 1990 Phase II study with an LH-RH analog in advanced ovarian cancer. *Journal of Gynecology and Endocrinology* **VI** 5–7.
- Schwartz PE, Keating G, MacLusky N, Naftolin F & Eisenfeld A 1982 Tamoxifen therapy for advanced ovarian cancer. *Obstetrics and Gynecology* **59** 583–588.
- Schwartz PE, Chambers JT, Kohorn EL, Chambers SK, Weitzman H, Voynick JM *et al.* 1989 Tamoxifen in combination with cytotoxic chemotherapy in advanced epithelial ovarian cancer. *Cancer* **63** 1074–1078.
- Sevela P, Denison U, Schemper M, Spona J, Varva N & Salzer H 1990 Oestrogen and progesterone receptor content as a prognostic factor in advanced epithelial ovarian carcinoma. *British Journal of Obstetrics and Gynecology* **97** 706–712.
- Shirley DR, Kavanagh JJ, Gershenson DM, Freedman RS, Copeland LJ & Jones LA 1985 Tamoxifen therapy for epithelial ovarian cancer. *Obstetrics and Gynecology* **66** 575–578.
- Sikic BI, Scudder SA, Ballon SC, Soriero OM, Christman JE, Suey L *et al.* 1986 High dose megestrol acetate therapy of ovarian carcinoma; a phase II study by the Northern California Oncology Group. *Seminars in Oncology* **4** 26–32.
- Simons WE, Albrecht M, Hansel M, Dietl M & Holzel F 1983 Cell line derived from human ovarian carcinomas: growth stimulation by gonadotropic and steroid hormones. *Journal of the National Cancer Institute* **70** 839–845.
- Simpson BJB, Rabiaz GJ, Macloed KG, Hirst GL, Miller WR & Langdon SP 1996 Estrogen regulation of TGF α in ovarian cancer. *Proceedings of the American Association for Cancer Research* **37** 224.
- Slevin ML, Harvey VJ, Osborne RJ, Shepherd JH, Williams CJ & Mead GM 1986 A phase II study of tamoxifen in ovarian cancer. *European Journal of Cancer and Clinical Oncology* **22** 309–312.
- Slotman BJ & Rao BR 1988 Ovarian cancer (review): etiology, diagnosis, prognosis, surgery, radiotherapy, chemotherapy and endocrine therapy. *Anticancer Research* **8** 417–434.

- Smith LH & Oi RH 1984 Detection of malignant ovarian neoplasms: a review of the literature. I Detection of the patient at risk; clinical, radiological and cytological detection. *Obstetrical and Gynecological Survey* **39** 313–328.
- Sutton GP, Senior MB & Strauss JF 1986 Estrogen- and gestagenen in epithelial ovarian malignancies. *Gynecology and Oncology* **23** 176–182.
- Thigpen JT, Vance RB, Balducci L & Khansur T 1984 New drugs and experimental approaches in ovarian cancer treatment. *Seminars in Oncology* **11** 314–326.
- Trope CG & Makar AP 1991 Epidemiology, etiology, screening, prevention, and diagnosis in female genital cancer. *Current Opinion in Oncology* **3** 909–919.
- Trope C, Johnsson JE, Sigurdsson K & Simonse E 1982 High-dose medroxyprogesterone acetate for the treatment of advanced ovarian carcinoma. *Cancer Treatment Reviews* **66** 1441–1443.
- Van Der Vange N, Gregg S, Burger W, Kenemans P & Vermorken JB 1995 Experience with hormonal therapy in advanced epithelial ovarian cancer. *Acta Oncologica* **34** 813–820.
- Van Der Velden J, Gitsch G, Wain GV, Friedlander ML & Hacker NF 1995 Tamoxifen in patients with advanced epithelial ovarian cancer. *International Journal of Gynecological Cancer* **5** 301–305.
- Van Leeuwen FE & Rookus MA 1989 The role of exogenous hormones in the epidemiology of breast, ovarian and endometrial cancer. *European Journal of Cancer and Clinical Oncology* **25** 1961–1972.
- Weiner SA, Alberts DS, Surwit EA, Davis J & Grosso D 1987 Tamoxifen therapy in recurrent epithelial ovarian carcinoma. *Gynecology and Oncology* **27** 208–213.
- Whittemore AS 1993 Fertility drugs and risk of ovarian cancer. *Human Reproduction* **8** 999–1000.
- Whittemore AS, Harris R & Itnyre J 1992 Characteristics relating to ovarian cancer risk: collaborative analysis of 12 US case–control studies. IV The pathogenesis of epithelial ovarian cancer. *American Journal of Epidemiology* **136** 1212–1220.
- Wilailak S, Linasmita V, Srisupundit S, Tangtrakul S, Bullangpoti S & Israngura N 1999 Phase II study of megestrol acetate in patients with refractory epithelial ovarian cancer. *International Journal of Gynaecological Cancer* **9** 12.
- Wu MI, Whittemore AS, Paffenbarger JR, Sarles DL, Kampert JB, Grosser S *et al.* 1988 Personal and environmental characteristics related to epithelial ovarian cancer in reproductive and menstrual events and oral contraceptive use. *American Journal of Epidemiology* **6** 1216–1227.
- Yano T, Pinski J, Halmos G, Szepeshazi K, Groot K & Schally AV 1994 Inhibition of growth of OV-1063 human epithelial ovarian cancer xenografts in nude mice by treatment with luteinizing hormone-releasing hormone antagonist SB-75. *Proceedings of the National Academy of Sciences of the USA* **91** 7090–7094.